


Adult Stage

After the adult mosquito emerges, it seeks a protective environment in the surrounding vegetation to allow its wings to complete development. Male mosquitoes tend to emerge prior to the female mosquito and will mate with the female as soon as she is capable. Female and male mosquitoes both require carbohydrate sources (nectar, plant exudates) throughout their life to maintain energy for flying, mating and seeking hosts for bloodmeals. Only the female mosquito takes a bloodmeal; she needs the extra protein to develop eggs. The process of taking a bloodmeal is how the mosquito is able to vector viruses, protozoans and helminthes (worms) to humans and animals. Male mosquitoes tend to live only a week or two while female mosquitoes can live for up to a month and produce multiple batches of eggs. Some mosquito species overwinter as blood fed females and can survive for multiple months.



Funding

Provided by the U.S. Environmental Protection Agency, Region VI office, Dallas, Texas.

Authors

Dr. Don L. Rennie, Extension Program Leader for Agricultural and Environmental Safety; Coordinator-Pesticide Safety Education Project (PSEP); Assistant Professor and Extension Specialist; and Dr. Mark Johnsen, Extension Assistant, Agricultural and Environmental Safety

June 2007



Mosquito Life Cycle

Mosquitoes are most often thought of as blood-thirsty pests that spread disease and ruin our outdoor activities. The majority of the mosquito's life cycle, however, is spent in the water, and it is only the adult stage that adversely affects human and animal health. This knowledge offers us points in their life cycle to initiate control before they can reach the adult stage. Each stage in the mosquito life cycle has diverse morphology (how they look), but they all follow a general lifecycle which is described in detail in this brochure.

Visit our Web site:
www.lsuagcenter.com

Louisiana State University Agricultural Center
William B. Richardson, Chancellor
Louisiana Agricultural Experiment Station
David J. Boethel, Vice Chancellor and Director
Louisiana Cooperative Extension Service
Paul D. Coreil, Vice Chancellor and Director


Pub. 3065 (10M) 05/08

Issued in furtherance of Cooperative Extension work, Act. of Congress of May 8 and June 30, 1914, in cooperation with the United States Department of Agriculture. The Louisiana Cooperative Extension Service provides equal opportunities in programs and employment.



Mosquito Lifecycle Stages

All mosquitoes go through four distinct stages (egg-larval-pupal-adult) in their life cycle, which can vary in length depending on temperature and food resources. In the summertime it takes mosquitoes 7-10 days to totally complete their life cycle from egg to adult.



Egg Stage

Mosquito species are broken into two categories based on where in the environment they lay their eggs.

Floodwater mosquito species lay their eggs above the water



level in areas that are prone to periodic flooding. These oviposition (egg laying) sites range from cattle hoof prints to empty soda cans. The eggs from these mosquito species can persist in the environment for 3 months to 2 years and is the stage used to survive the winter.


Standing water mosquito species lay their eggs on the surface of transient and permanent pools of water. Some species lay their eggs singly while others glue 100-200 eggs together forming miniature rafts. Within 24-48 hours these eggs will hatch in the environment.

Larval Stage

Once hatched, mosquito larvae (or wigglers) go through 4 instars (time period between molts) increasing in size after each molt. The larval stage displays a great variation in morphology between the mosquito species and is used for identification purposes.



Most mosquito species feed by filtering micro-



organisms (bacteria, protozoa, fungi) out of the water, but several predacious species feed on other mosquito larvae. Larvae must breathe atmospheric oxygen and will lay horizontal (only *Anopheles* spp.) or 45 degrees to the water surface. Some species have the capability of puncturing underwater plant cells to receive oxygen and never have to surface.

Pupal Stage

The fourth instar larvae molts into the comma shaped stage called the pupa (or tumblers). This is the transition stage between the aquatic stages of the mosquito's lifecycle and the terrestrial adult stage. Mosquito pupae do not feed but are mobile and use a tumbling motion to escape predation. Within 24-48 hours the pupa will moult into an adult.

