

HOME GARDEN WEED MANAGEMENT**HOME GARDENS**

Active Ingredient	Product Rate per 1000 sq ft/1 gal¹	Remarks and Precautions
BURNDOWN:		
glyphosate ¹	Roundup Ultra/others @ 0.5-1 oz/1,000 sq ft. Several brands available; consult labels for proper rates.	Apply post-emergence on weeds. A nonselective post-emergent to be used before planting or crop emergence on actively growing weeds. Consult product label for plant back intervals. Avoid spray drift to desirable plants. Spray weeds to wet but not to drip. Crops labeled include beets, carrots, celery, corn, cole crops, English and Southern peas, Irish potatoes, Jerusalem artichoke, lima and snap beans, lettuce, greens, okra, onion, garlic, peanuts, radish, soybeans, sweet potatoes and asparagus (before spears appear).
PRE-PLANT INCORPORATED:		
trifluralin	Treflan 4 EC @ 1/3 fl oz/1,000 sq ft on medium soil 1/2 fl oz/1,000 sq ft on heavy soils	Apply before planting and incorporate. Use on snap or lima beans, cole crops, Irish potatoes, sunflowers and greens. Incorporate before transplanting tomatoes, cole crops, pepper and celery. Also for asparagus beds and at half-rate post-emergence (crop) between onion rows. Treflan controls many small-seeded broadleaf weeds (pigweeds) and annual grasses.
PRE-EMERGENCE (before crops or weeds emerge):		
metolachlor	Dual 8E @ 1/2 oz/1,000 sq ft on sandy soil 1.0 fl. oz/1,000 sq ft on medium-heavy soil	Apply before or after planting snap or lima beans, soybeans, sweet corn, peanuts, popcorn, Irish potatoes, Southern peas, English peas and chickpeas. Apply after planting snap or lima beans, sweet corn, peanuts, popcorn, Irish potatoes, southern pecans, English peas and chickpeas. Provides good residual grass control and controls some broadleaf weeds. Will not control emerged weeds.
DCPA	Dacthal 75% WP @ 4.0 oz/1,000 sq ft on sandy soil 5.0 oz./1,000 sq. ft. on med-heavy soil	Apply after planting snap beans (not limas), Southern peas, greens, radish, garlic, onion, cole crops, Irish or sweet potatoes. Apply after transplanting tomatoes, peppers, cole crops and cucurbits. Apply 4-6 weeks after transplanting eggplants. Controls many annual grasses and some broadleaf weeds.
POST-EMERGENCE (after crop and weeds emerge):		
bentazon	Basagran 4 EC @ 1/2-3/4 oz/1,000 sq ft	Snap or lima beans, mint, Southern peas, peanuts, English peas and corn. Apply to small actively growing broadleaf weeds and yellow nutsedge. This treatment will not kill grasses. Very effective on yellow nutsedge. May need an adjuvant; see label.
fluazifop	Fusilade DX @ 0.5 oz/1,000 sq ft Apply with NIS @ 0.33 oz/gal	Spray on young actively growing grassy weeds. Spray to wet weed foliage. Use on asparagus, carrots, onions, spinach, sweet potatoes, hot peppers and non-bell peppers.

HOME GARDEN WEED MANAGEMENT**HOME GARDENS**

Active Ingredient	Product Rate per 1000 sq ft/1 gal¹	Remarks and Precautions
POST-EMERGENCE (after crop and weeds emerge) continued:		
sethoxydim	Poast & various other trade names @ 1 oz/1000 sq ft Consult product label concerning surfactants	Spray on young, actively growing grassy weeds. Spray to wet weed foliage. Use on asparagus, beans, cole crops, cabbage, cucumbers, peanuts, greens, lettuce, melons, peas, peppers, pumpkins, potatoes, squash or tomatoes. Sethoxydim may be found under various trade names like Poast, Hi-Yield Grass Killer and Ferti-lome Over the Top II.

¹ The rates given in column 2 (Product Rate) are based on the amount of material needed to treat 1,000 square feet. This amount of herbicide plus adjuvant (if required) should be diluted in 1 gal. of water and evenly distributed over 1,000 sq. ft. To find the amount of product needed if the area treated is different from 1,000 sq. ft., use the following formula: [product rate, in oz. (column 2) × (area to be treated, in sq. ft. ÷ 1,000)]. To find the volume necessary to spray, use the following formula: [area to be treated, in sq. ft. ÷ 1,000]. Example: area to be treated is 2,500 sq. ft. Want to spray Fusilade DX @ 0.5 oz/1,000 sq. ft. Total product needed: [0.5 × (2,500 ÷ 1000)] or 1.25 oz. Total volume needed is [2,500 ÷ 1,000] or 2.5 gal.

² Refer to "Appendix A" for more information on various glyphosate brands and formulations.