



Your Experts for Life

ANR-174

# Nutrient Content of Fertilizer Materials

The following tables can be used as quick reference guides to fertilizer materials. These materials may be used alone or blended with other fertilizer materials to form a multinutrient fertilizer.

The actual nutrient content may vary from what is listed, depending upon the manufacturer, the

purity of the product, or other materials blended with the product. Most values are for the fertilizer-grade product and not the pure chemical. The chemical formulas given are for the primary active compound.

## Primary and Secondary Nutrient Sources

Material	Percentage						Approx. CaCO <sub>3</sub> Equiv./100 lb.†	Comments
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	Ca	Mg	S		
<b>Nitrogen Materials</b>								
<b>Ammonium nitrate‡</b> NH <sub>4</sub> NO <sub>3</sub>	34	0	0	0	0	0	-61	
<b>Ammonium nitrate limestone</b> NH <sub>4</sub> NO <sub>3</sub> + (CaCO <sub>3</sub> + MgCO <sub>3</sub> )	20	0	0	6	4	0	0	Ca and Mg depend on limestone used
<b>Ammonium nitrate sulfate</b> NH <sub>4</sub> NO <sub>3</sub> + (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	30	0	0	0	0	5	-71	
<b>Ammonium sulfate‡</b> (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	21	0	0	0	0	24	-110	
<b>Anhydrous ammonia‡</b> NH <sub>3</sub>	82	0	0	0	0	0	-148	Pressurized gas
<b>Aqua ammonia</b> NH <sub>4</sub> OH	16-25	0	0	0	0	0	-36 to -54	
<b>Calcium cyanamide</b> CaCN <sub>2</sub>	21	0	0	11	0	0	+63	Most alkaline N material
<b>Calcium nitrate</b> Ca(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	15	0	0	21	0	0	+20	
<b>Calcium nitrate/urea (Calurea)</b> Ca(NO <sub>3</sub> ) <sub>2</sub> + 4CO(NH <sub>2</sub> ) <sub>2</sub>	34	0	0	10	0	0	-36	Don't blend with superphosphate
<b>Crotonylidene diurea</b> (CDU)	32	0	0	0	0	0	NA	Slow-release
<b>Isobutylidene diurea</b> (IBDU)	31	0	0	0	0	0	NA	Slow-release
<b>Nitrogen solutions (N-SOL or UAN solutions)‡</b> (urea/ammonium nitrate):								Solutions code:
32% UAN (35% urea + 45% A.N.)	32	0	0	0	0	0	-55	320(0-45-35)
30% UAN (33% urea + 42% A.N.)	30	0	0	0	0	0	-52	300(0-42-33)
28% UAN (30% urea + 40% A.N.)	28	0	0	0	0	0	-49	280(0-40-30)
21% AN (60% A.N. + 40% water)	21	0	0	0	0	0	-37	210(0-60-0)
19% AN (54% A.N. + 46% water)	19	0	0	0	0	0	-33	190(0-54-0)
<b>Potassium nitrate</b> KNO <sub>3</sub>	13	0	44	0	0	0	+26	
<b>Sodium nitrate</b> (nitrate of soda) NaNO <sub>3</sub>	16	0	0	0	0	0	+29	
<b>Urea‡</b> CO(NH <sub>2</sub> ) <sub>2</sub>	45	0	0	0	0	0	-81	
<b>Urea</b> (sulfur coated) CO(NH <sub>2</sub> ) <sub>2</sub> +S	36-38	0	0	0	0	13-16	-118	N release depends on S coating
<b>Ureaform</b> (urea + formaldehyde)	38	0	0	0	0	0	-68	Slow-release

Material	Percentage						Approx. CaCO <sub>3</sub> Equiv./100 lb.†	Comments
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	Ca	Mg	S		
<b>Phosphorus Materials</b>								
<b>Ammoniated superphosphate</b>	12-17	22-35	0	*	0	*	-7	
<b>Ammonium phosphate‡</b>								
Diammonium phosphate (DAP) (NH <sub>4</sub> ) <sub>2</sub> H <sub>2</sub> PO <sub>4</sub>	18	46	0	0	0	0	-70	
Monoammonium phosphate (MAP) NH <sub>4</sub> HPO <sub>4</sub>	11	48	0	1	0	0	-65	
<b>Ammonium phosphate nitrate</b> NH <sub>4</sub> H <sub>2</sub> PO <sub>4</sub> ·NH <sub>4</sub> NO <sub>3</sub>	30	10	0	0	0	0	-54	
<b>Ammonium phosphate sulfate</b> 4NH <sub>4</sub> H <sub>2</sub> PO <sub>4</sub> ·x(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	16	20	0	0	0	15	-76 to -113	Mixture
<b>Ammonium polyphosphate‡</b> (NH <sub>4</sub> )HP <sub>2</sub> O <sub>7</sub> + (NH <sub>4</sub> ) <sub>5</sub> P <sub>3</sub> O <sub>10</sub>	10	34	0	0	0	0	—	Liquid
<b>Basic Slag</b>	0	0-6	*	3-29	*	*	+70	Analysis variable
<b>Bone meal</b> (steamed) Ca <sub>3</sub> (PO <sub>4</sub> ) <sub>2</sub> + CaCO <sub>3</sub>	0-2	10-20	0	19-25	0	0	+20	
<b>Concentrated superphosphate‡</b> (triple superphosphate) Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub> ·xH <sub>2</sub> O	0	46	0	14	0	2	0	
<b>Nitric phosphate</b>	12-17	22-35	0	*	0	*	-20	
<b>Normal superphosphate</b> Ca(H <sub>2</sub> PO <sub>4</sub> ) <sub>2</sub> ·xH <sub>2</sub> O + CaSO <sub>4</sub>	0	20	0	21	0	11	0	
<b>Phosphate rock</b>	0	2-35	0	*	*	0	+10	Total P <sub>2</sub> O <sub>5</sub> relatively unavailable
<b>Phosphoric acid‡</b> H <sub>3</sub> PO <sub>4</sub>								Liquid
Wet-process acid	0	30	0	0	0	0	-63	
Concentrated wet-process acid	0	40-54	0	0	0	0	-84 to -113	
Superphosphoric acid	0	76	0	0	0	0	-110	
<b>Urea-ammonium phosphate</b>	25	35	0	0	0	0		Research mixture
<b>Urea phosphate</b> CO(NH <sub>2</sub> ) <sub>2</sub> + H <sub>3</sub> PO <sub>4</sub>	17	44	0	0	0	0	-82	Research
<b>Potassium Materials</b>								
<b>Greensand</b>	0	1	6	0	0	0		Natural low-grade, mineral
<b>Potassium carbonate</b>								
K <sub>2</sub> CO <sub>3</sub> solid	0	0	48	0	0	0	+70	
K <sub>2</sub> CO <sub>3</sub> liquid	0	0	34	0	0	0	+50	
<b>Potassium chloride‡</b> KCl(muriate of potash)	0	0	60	0	0	0	0	Most widely used, single fertilizer material
<b>Potassium magnesium sulfate‡</b> (sulfate of potash magnesia) K <sub>2</sub> SO <sub>4</sub> ·2MgSO <sub>4</sub> or MgSO <sub>4</sub> ·K <sub>2</sub> SO <sub>4</sub> ·6H <sub>2</sub> O	0	0	21	0	11	23	0	
<b>Potassium metaphosphate</b> KPO <sub>3</sub>	0	59	39	0	0	0	*	
<b>Potassium nitrate</b> (nitrate of potash) KNO <sub>3</sub>	13	0	44	0	0	0	+26	
<b>Potassium sulfate</b> (sulfate of potash) K <sub>2</sub> SO <sub>4</sub>	0	0	52	0	0	16	0	
<b>Calcium Materials</b>								
<b>Calcium chloride</b> CaCl <sub>2</sub>	0	0	0	36	0	0	0	Water soluble
<b>Burned lime</b> CaO	0	0	0	70	0	0	+178	
<b>Calcitic limestone‡</b> (ground) CaCO <sub>3</sub>	0	0	0	36	0	0	+95 to 100	
<b>Dolomitic limestone‡</b> (ground) CaCO <sub>3</sub> + MgCO <sub>3</sub>	0	0	0	24-30	6-12	0	+95 to 108	
<b>Selma chalk</b>	0	0	0	32	0	0	+80	
<b>Gypsum‡</b> CaSO <sub>4</sub> ·2H <sub>2</sub> O	0	0	0	22	0	18	0	Solubility = 0.02 lb./gal.
<b>Hydrated lime</b> Ca(OH) <sub>2</sub>	0	0	0	50	0	0	+134	Solubility = 0.01 lb./gal.

Material	Percentage						Approx. CaCO <sub>3</sub> Equiv./100 lb.†	Comments
	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	Ca	Mg	S		
<b>Magnesium Materials</b>								
<b>Dolomitic limestone‡</b> (ground) CaCO <sub>3</sub> + MgCO <sub>3</sub>	0	0	0	24-30	6-12	0	+95 to +108	Analysis variable; must contain at least 6% Mg
<b>Magnesium ammonium phosphate</b> MgNH <sub>4</sub> PO <sub>4</sub> ·6H <sub>2</sub> O	8	40	0	0	15	0		Solubility = 0.001 lb./gal.
<b>Magnesium oxide</b> MgO	0	0	0	0	45	0	+250	Solubility = 0.00062 g/100 g 0.001 oz./gal.
<b>Magnesium sulfate‡</b> (Epsom salt) MgSO <sub>4</sub> ·7H <sub>2</sub> O	0	0	0	0	10	13	0	Solubility = 7.6 lb./gal.
<b>Magnesium sulfate</b> (Kieserite) MgSO <sub>4</sub> ·H <sub>2</sub> O	0	0	0	0	17	23	0	Solubility = 5.7 lb./gal.
<b>Potassium magnesium sulfate‡</b> (sulfate of potash magnesia) K <sub>2</sub> SO <sub>4</sub> ·2MgSO <sub>4</sub>	0	0	21	0	11	23	0	Soluble
<b>Sulfur Materials</b>								
<b>Ammonium sulfate‡</b> (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	21	0	0	0	0	24	-110	Available in prilled or water-soluble crystals
<b>Ammonium thiosulfate‡</b> (60% solution) (NH <sub>4</sub> ) <sub>2</sub> S <sub>2</sub> O <sub>3</sub>	12	0	0	0	0	26		Liquid; reacts with alkaline materials
<b>Elemental sulfur (S):</b>								Elemental S must be oxidized to sulfate before available to plants
Wettable S	0	0	0	0	0	90-100	-312	
Flowable S	0	0	0	0	0	52-70	-218	
Flowers of S	0	0	0	0	0	90-100	-312	
<b>Gypsum‡</b> CaSO <sub>4</sub> ·2H <sub>2</sub> O	0	0	0	22	0	18	0	
<b>Magnesium sulfate</b> (Epsom salt) MgSO <sub>4</sub> ·7H <sub>2</sub> O	0	0	0	0	10	13	0	Soluble
<b>Potassium magnesium sulfate‡</b>	0	0	21	0	11	23	0	
<b>Potassium sulfate</b> K <sub>2</sub> SO <sub>4</sub>	0	0	52	0	0	16	0	Soluble
<b>Sulfuric acid</b> H <sub>2</sub> SO <sub>4</sub>	0	0	0	0	0	20-26	-62 to -81	Liquid; highly reactive

†Negative value indicates net acidifying effect on soil; positive value indicates net basic reaction in soil.

‡Commonly available materials

\*Present in undetermined amounts

Symbol key: N = Nitrogen; P<sub>2</sub>O<sub>5</sub> = Phosphate; K<sub>2</sub>O = Potash; Ca = Calcium; Mg = Magnesium; S = Sulfur; CaCO<sub>3</sub> = Calcitic Limestone

### Micronutrient Materials

Materials	Nutrient Content	Materials	Nutrient Content
<b>Copper (Cu)</b>		<b>Boron (B)</b>	
Chelated Cu*:		Solubor*	20% B
Cu EDTA	13% Cu	Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub>	66% B <sub>2</sub> O <sub>3</sub>
Cu HEDTA	9% Cu	Magnesium borate (boracite)	
Cupric ammonium phosphate		2Mg <sub>3</sub> B <sub>8</sub> O <sub>15</sub> ·MgCl <sub>2</sub>	21% B
Cu(NH <sub>4</sub> )PO <sub>4</sub> ·H <sub>2</sub> O	30% Cu	<b>Iron (Fe)</b>	
Cupric oxide (CuO)	60-80% Cu	Basic slag	10-13% Fe
Copper sulfate*:		Ferric sulfate Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> ·4H <sub>2</sub> O	20% Fe
CuSO <sub>4</sub> ·H <sub>2</sub> O	35% Cu	Ferrous sulfate* FeSO <sub>4</sub> ·7H <sub>2</sub> O	20% Fe
CuSO <sub>4</sub> ·5H <sub>2</sub> O	25% Cu	Ferrous ammonium sulfate	
CuSO <sub>4</sub> ·3Cu(OH) <sub>2</sub>	13-53% Cu	(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> ·FeSO <sub>4</sub> ·6H <sub>2</sub> O	14% Fe
Copper frits	40-50% Cu	Ferrous ammonium phosphate	29% Fe
Copper polyflavonoid	6% Cu	Fe(NH <sub>4</sub> )PO <sub>4</sub> ·H <sub>2</sub> O	
<b>Boron (B)</b>		Ferrous oxalate FeC <sub>2</sub> O <sub>4</sub> ·2H <sub>2</sub> O	30% Fe
Borax (sodium tetraborate decahydrate)		Ferrous carbonate	
Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub> ·10H <sub>2</sub> O	11% B	FeCO <sub>3</sub> ·H <sub>2</sub> O	42% Fe
Boric acid (H <sub>3</sub> BO <sub>3</sub> )	17% B	Iron chelates*:	
Boron frit/sodium borosilicate	6% B	Fe DTPA	10% Fe
Calcium borate (colemanite)		Fe EDTA	9-12% Fe
Ca <sub>2</sub> B <sub>6</sub> O <sub>11</sub> ·5H <sub>2</sub> O	10% B	Fe EDDHA	6% Fe
Fertilizer borate* (sodium tetraborate)		Fe HEDTA	5-9% Fe
Borate Granular (Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub> ·5H <sub>2</sub> O)	14% B	Iron ligninsulfonate	5-11% Fe
Borate 48	15% B	Iron polyflavonoid	6-10% Fe
	48 refers to percentage B <sub>2</sub> O <sub>3</sub>	Iron Frits	40% Fe

Materials	Nutrient Content
<b>Manganese (Mn)</b>	
Basic slag	1-3% Mn
Manganese frits	10-25% Mn
Manganese chloride MnCl <sub>2</sub>	17% Mn
Manganese carbonate MnCO <sub>3</sub>	31% Mn
Manganese oxide MnO	68-70% Mn
Manganese sulfate* MnSO <sub>4</sub> ·4H <sub>2</sub> O	24% Mn
Manganese chelate* Mn EDTA	12% Mn
Manganese ammonium phosphate Mn(NH <sub>4</sub> )PO <sub>4</sub> ·H <sub>2</sub> O	28% Mn
Manganese polyflavonoid	8% Mn
<b>Molybdenum (Mo)</b>	
Ammonium molybdate (NH <sub>4</sub> ) <sub>6</sub> Mo <sub>7</sub> O <sub>24</sub> ·2H <sub>2</sub> O	up to 54% Mo

Materials	Nutrient Content
<b>Molybdenum (Mo)</b>	
Sodium molybdate* Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	38-46% Mo
Molybdenum frit	30% Mo
Superphosphate	trace as impurity
<b>Zinc (Zn)</b>	
Zinc chelate* Na <sub>2</sub> Zn EDTA	9-14% Zn
Zinc ammonium phosphate Zn(NH <sub>4</sub> )PO <sub>4</sub> ·H <sub>2</sub> O	34% Zn
Zinc sulfate* ZnSO <sub>4</sub> ·H <sub>2</sub> O	22-36% Zn
Zinc sulfide (sphalerite)	61% Zn
Zinc oxide ZnO	78-80% Zn
Zinc ligninsulfonate	5-12% Zn
Zinc polyflavonoid	7-10% Zn

\*Commonly used materials

### Organic Fertilizer Materials (Approximate Values)

Material	N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O	Ca	Mg	S	Micro-nutrients	Acidic (-)		Comments
								Neutral (0)	Alkaline (+)	
<b>Blood</b> (dried)	12-15	3	1	*	*	*	*	-		
<b>Blood meal</b>	15	1	1	*	*	*	*	-		
<b>Bone meal</b> (steamed)	0-2	10-20	0	19-25	0	0		+		
<b>Compost</b> (garden)	Variable depending upon components and amendments									
<b>Cottonseed meal</b>	6-7	2.5	1.5	*	*	*	*	-		
<b>Cottonseed hull ash</b>	0		27	*	*			+		
<b>Cotton notes</b> (composted gin wastes)	2	0.5	3	4	0.7	0.6	*	-		May contain weed seed
<b>Fish scrap</b> (acidulated)	7-10	1-2	0	*	*	2		-		Traces of I and Hg
<b>Fish scrap</b> (dried fish meal)	9	3		6	*	*	*			
<b>Hay:</b>										
Legume	3.0	1.0	2.4	1.2	0.2	0.3	*	-		
Grass	1.5	0.5	1.9	0.8	0.2	0.2	*	-		
<b>Manure:</b> (dried)										
Cattle	1.5	1.5	1.2	1.1	0.3	*	*	-		Feedlot manure usually contains 60-70% water
Horse	0.4	0.2	0.3	*	*	*	*	-		
Poultry										
Broiler litter	3.0	3.0	2.0	1.8	0.4	0.3	*	+		Average moisture (houses) = 20%
Hen-caged layers	1.5	1.3	0.5	6	0.4	0.3	*	+		Average moisture = 70%
Hen-litter	1.8	2.8	1.4	*	*	*	*	+		Average moisture = 37%
Sheep	0.6	0.3	0.2	*	*	*	*	-		
Swine	0.6	0.4	0.1	*	*	*	*	-		
<b>Peat/Muck</b>	2.3	0.5	0.7		*	*				
<b>Sawdust</b> (mixed soft + hardwoods)	0.2	0	0.2					-		High C:N ratio; needs N fert.
<b>Seaweed</b> (dried)	0.7	0.8	5.0	*	*	*	*	+, -		
<b>Sewage sludge</b> (dried, municipal)	5	6	0.5	3	1	1		+		Depending upon source, may contain heavy metals. Use only on nonfood-producing areas.
<b>Tankage</b>	7	1.5	3-10	*	*	*	*	-, +		
<b>Wood ashes</b>	0	2	6	20	1		*	+		70% CaCO <sub>3</sub> equivalent

\*Present in undetermined amounts      Symbol key: N = Nitrogen; P<sub>2</sub>O<sub>5</sub> = Phosphorus; K<sub>2</sub>O = Potassium; Ca = Calcium; Mg = Magnesium; S = Sulfur



**Charles C. Mitchell**, *Extension Agronomist*, Professor, Agronomy and Soils, Auburn University

**For more information**, call your county Extension office. Look in your telephone directory under your county's name to find the number.

Issued in furtherance of Cooperative Extension work in agriculture and home economics, Acts of May 8 and June 30, 1914, and other related acts, in cooperation with the U.S. Department of Agriculture. The Alabama Cooperative Extension System (Alabama A&M University and Auburn University) offers educational programs, materials, and equal opportunity employment to all people without regard to race, color, national origin, religion, sex, age, veteran status, or disability.

6M, Reprinted March 2008, ANR-174