

# Fruit and Nuts – Commercial

## Fig Spray Schedule

Insect or Arthropod	Material	Rate	Restrictions
Vinegar flies	Malathion 57 EC (Malathion)	2 quarts plus 1-2 gallons unsulfurized molasses/acre	PHI=3 days; use 300 gallons water/acre. <u>Sanitation:</u> Early harvest and complete fruit removal at harvesting will reduce fruit exposure to flies. Disc under affected hosts to destroy fermented fruit residues.
Fig scale	Volk supreme oil	3 gallons/100 gallons water	Dormant or delayed dormant spray only.
Spider mites	Omite 30WP (Propargite)	6 pounds/acre Two applications	Use only on nonbearing trees and when trees will not bear fruits within 1 year of application.
	Omite 6E (Propargite)	2 pints/acre Two applications	
	Volk supreme oil	3 gallons/100 gallons water	Dormant or delayed dormant spray only.
	Sulfur	Follow labeled rate	Do not mix with oils and/or do not apply within 3 weeks of oil applications to avoid fruit burn. Some sulfur formulations should not be combined with spreader stickers. Sulfur may cause eye and skin irritation. Avoid when hot temperatures (above 90 F) are expected within three days of spraying.

**Warning:** Re-entry time for workers entering treated areas should be strictly observed. Read the label for this information.