

# Fruit and Nuts – Commercial

## Mayhaw Spray Schedule

Insect	Material	Rate	Restrictions
<b>Aphids</b> (except wooly apple aphid)	Admire Pro 4F (Imidacloprid)	2.8 fluid ounces/per acre	PHI=7 days; apply postbloom; highly toxic to bees.
	Mustang Max (Zeta-cyprmethrin)	0.21-0.32 fluid ounce/50 gallons or 1.28 fluid ounces/acre	PHI=14 days; allow 7 days between applications.
<b>Leafminers</b>	SpinTor 2SC (Spinosad)	1.0-2.5 fluid ounces/100 gallons or 4.0-10.0 fluid ounces/acre	PHI=7 days; do not apply more than 29 fluid ounces/acre/year.
<b>Leafhoppers</b>	Admire Pro 4F (Imidacloprid)	1.4 – 2.8 fluid ounces/per acre	PHI=7 days; apply post-bloom; highly toxic to bees.
	Mustang Max (Zeta-cyprmethrin)	0.21-0.32 fluid ounce/50 gallons or 1.28 fluid ounces/acre	PHI=14 days; allow 7 days between applications.
<b>Aphids, plum curculio</b>	Actara (Thiamethoxam)	4.5 ounces/acre	PHI=35 days; highly toxic to bees; do not make more than 1 prebloom application.
	Mustang Max (Zeta-cyprmethrin)	0.21-0.32 fluid ounce/50 gallons or 1.28 fluid ounces/acre	PHI=14 days; allow 7 days between applications.
<b>Spider mites</b>	Savey 50DF (Hexythiazox)	3.0-6.0 ounces/acre	PHI=28 days; apply at the beginning of mite infestations. Savey is not effective against adult mites. Limit to one application/year. Do not use in home plantings.

**Warning:** Re-entry time for workers entering treated areas should be strictly observed. Read the label for this information.