

Crops - Commercial

Wheat and Oats					
Insect	Insecticide	Amount of Concentrate per Acre	Pounds Active Ingredient/ Acre	Acres Treated per Gallon or Pound SP	Comments
Aphids (including greenbug aphid)	malathion Malathion (5)	24 ounces	1	5.3	General guideline for greenbug treatment levels:
	sulfoxaflor Transform (50)	0.75 ounce	0.023	21.3	Plant height (inches): 3-6, 4-8, 6-16 Number of greenbugs (per linear foot): 100-300, 200-400, 300-800
	dimethoate Dimethoate (4)	8-12 ounces	0.25-0.375	16-10.7	Wheat only.
	SUPPRESSION				
	lambda-cyhalothrin Karate Z (2.08)	1.92 ounces	0.03	66.7	Wheat only. See labels. Treat when greenbugs cause areas with dead plants. Other aphid species may not require control.
	z-cypermethrin Mustang Max (0.8)	3.2-4 ounces	0.02-0.025	40-32	
True armyworm, fall armyworm	lambda-cyhalothrin Karate Z (2.08)	1.28-1.92 ounces	0.02-0.03	100-66.7	Treat when 5 or more worms per square foot are found, and foliage loss is occurring.
	spinosad Blackhawk (0.36)	1.7-3.3 ounces	0.038-0.075	9.4-4.8	
	z-cypermethrin Mustang Max (0.8)	3.2-4 ounces	0.02-0.025	40-32	Wheat only.
Stink bugs	z-cypermethrin Mustang Maxx (0.8)	3.2-4 ounces	0.02-0.025	40-32	Wheat only.
	lambda-cyhalothrin Karate Z (2.08)	1.28-1.92 ounces	0.02-0.03	100-66.7	Treat when 10% of spikes (wheat heads) in the milk stage are infested. Treat when 25% of spikes in the soft dough stage are infested.
Hessian fly	Host plant resistance and cultural control tactics				
	<ol style="list-style-type: none"> 1. Plant resistant varieties where available. Use of resistant varieties is the key method for management of Hessian fly. The resistant variety must have the resistant gene(s) for the Hessian fly biotype present. 2. Plant during recommended time period. There is no "fly-free date" for Louisiana. 3. Rotate wheat ground, if possible. 4. Use conventional tillage in the spring. 5. Eliminate volunteer wheat in the summer (following harvest) and fall (prior to harvest). 6. Avoid using susceptible wheat for cover crops and wildlife plantings. 				

Crops - Commercial

Wheat and Oats					
Insect	Insecticide	Amount of Concentrate per Acre	Pounds Active Ingredient/ Acre	Acres Treated per Gallon or Pound SP	Comments
Hessian fly - Continued	Seed treatments				
	imidacloprid Gaucho 600	1.6 fluid ounces per cwt.			Wheat only.
	imidacloprid Gaucho XT	3.4 fluid ounces + Gaucho 600 @ 1 fluid ounce/cwt			Wheat only.
	thiamethoxam Cruiser 5FS	1.33 fluid ounce per cwt.			Wheat only.
	thiamethoxam Cruiser Maxx Cereals	5 fluid ounces + Cruiser 5S @ 0.5 – 1 fluid ounce/cwt.			Wheat only.
	clothianidin Nipsit INSIDE	1.79 fluid ounce per cwt			Wheat only
	Foliar treatments				
	lambda-cyhalothrin Karate Z (2.08)	1.92 ounce	0.03	66.7	Wheat only. Apply when adult Hessian fly are emerging and active in the fall (2- to 4-leaf stage) or the late winter, early to mid-March). Timing of applications is difficult.

Insecticide Precautions and Limitations (refer to insecticide label for complete information)
Malathion: Do not apply within 7 days of harvest. REI: 12 hours.
Dimethoate: Do not apply within 14 days of grazing or 35 days of harvest. REI: 48 hours.
Karate Z: Do not apply within 30 days of harvest. Waiting period for grazing is 7 days. Do not apply more than 0.06 pound per acre per season. REI: 24 hours.
Tracer: Preharvest treatment interval: 21 days for grain, 14 days for forage or hay harvest. Do not apply more than 0.28 pound AI per acre per year. REI: 4 hours.
Mustang Maxx: Do not apply more than 0.125 pound AI per acre per season. Postharvest interval is 14 days. REI: 12 hours.
Cruiser: Toxic to wildlife and aquatic organisms. REI: 12 hours.
Gauchos: Toxic to birds and aquatic invertebrates. Do not graze livestock for 45 days. REI: 12 hours.