

2020-2021



Cool-Season



Pasture & Forage Varieties

Variety Selection

One of the most important decisions a forage producer must make is which variety or varieties to plant. Many varieties of forage crops are marketed in Louisiana. To help farmers make logical choices among crops and varieties, the LSU AgCenter conducts variety trials and makes suggestions each year. These suggestions are available in your county agent's office (Table 1).

Promising Varieties: Promising varieties are those that have been tested on a limited basis for two years and found to perform well enough to suggest trial plantings. These varieties are not yet fully suggested, and producers should avoid large plantings until further testing can be completed.

Grass Varieties

Fescue: Fescue is better adapted to heavier soils of northern Louisiana than to other soils or locations in the state. In recent years the variety Jesup Max Q tall fescue has been released and is available as certified low endophyte seed. This variety yields well but appears to be less persistent than older varieties, such as Kentucky-31. For maximum survival, these varieties should be well established before grazing begins and should not be grazed below a 2-inch height. No nitrogen should be applied from May 1 to September. Harvesting fescue for hay is not advised during the period from June 1 to September.

Oats: Oats can be successfully planted slightly earlier and produce earlier growth than ryegrass or wheat. They are often included in grazing mixtures for these reasons. Varieties of oats are tested and suggested primarily on the basis of dry forage yields. The yield ranking among suggested varieties has varied from location to location and year to year. In recent years RAM LA 99016 has been among the leaders (Table 2).

Rye: Rye is a good producer of fall, winter and early spring growth.

It matures earlier than ryegrass or other small grains. Rye is sometimes planted to give early grazing. Varieties of rye are tested and suggested primarily on the basis of dry forage yields. Rye varieties have not been evaluated in recent years.

Ryegrass: Ryegrass is important as a cool-season annual grass for grazing and harvested forage in Louisiana. Its high yielding ability, ease of establishment, high forage quality and tendency to form a denser sod than small grains make it attractive. Its peak growth period is in the spring, but it has good fall growth if planted early and weather is suitable.

Varieties of ryegrass are tested annually at several locations.

In addition to yield determination and seasonal yield distribution at each location, rust ratings have been made at the Iberia Research Station at Jeanerette when conditions permitted. In those evaluations, Jackson has generally been least infected by rust. Marshall and Gulf have been most severely infected.

Yield rankings of ryegrass varieties have varied with year and location (Table 3). Gulf has exhibited poor cold tolerance, and its yield performance has been inconsistent.

Wheat: Wheat is a good producer of late fall and spring growth.

The LSU AgCenter only does a limited amount of testing of wheat varieties for forage. No specific variety suggestions are presented because of a lack of data.

Triticale: Triticale is a genetic cross between wheat and rye. It can be grown for grain or forage. It is more winter hardy than oats, but similar to wheat and rye. The variety FL 08128 is the only variety suggested for planting.

Legume Varieties

Alfalfa: Alfalfa is an important hay and haylage crop in many states, but the amount produced in Louisiana is very limited.

It requires excellent drainage, highly fertile soils, a near neutral soil pH and a high level of management. Many varieties are marketed, but only a few are adapted in Louisiana.

Arrowleaf Clover: Arrowleaf clover is an annual with fair reseeding ability. It starts growth in the fall but produces most of its growth from March to late May or even to July, depending on moisture supply and variety. Varieties suggested in Louisiana

include Amclo (early maturity), Yuchi (intermediate maturity) and Meechi (late maturity). The difference in maturity between Amclo and Meechi is about five weeks.

Berseem Clover: Berseem clover is an annual, upright-growing clover. The only variety suggested in Louisiana is Bigbee, which was first suggested in 1986. It is noted for excellent fall and winter growth and a long period of good growth in the spring.

Its reseeding ability in Louisiana has been disappointing, but it has produced reseeding stands in Mississippi.

Crimson Clover: Crimson clover is an upright-growing annual clover. It produces some fall and winter growth but produces most of its growth in early spring. All suggested varieties are reseeding varieties. Dixie and Tibbee are early maturing, and Chief is late maturing.

Red Clover: Red clover is an upright-growing clover that is a short-lived perennial. In Louisiana, it is managed as an annual or biennial. It has poor reseeding ability and depends on most of its growth in late spring or early summer. There are several new varieties available that appear to be promising.

Subterranean Clover: Subterranean clover is a low-growing annual clover that has prostrate creeping stems with erect leaves. Seeds are produced in a bur that develops at or below the soil surface. It has excellent reseeding ability under close grazing, but some reseeding failures have been reported after the initial stand had thrived for several years. It produces most of its growth in the spring, slightly later than crimson clover.

All of the suggested varieties originated in Australia. In Australia, the maturity is considered to be early for Woogenellup, midseason for Mt. Barker, and late for Nangeela and Tallarook. Woogenellup has a low to moderate level of hard-seededness. The others have a low level of hard-seededness.

White Clover: White clover is a low-growing, perennial clover that has prostrate stems and erect leaves. A few varieties have good reseeding ability and act more like annuals than perennials in Louisiana. White clover produces most of its growth in the spring, early summer and late fall.

Louisiana S-1 is an intermediate variety that has intermediate size and good reseeding ability. It has some perennial tendencies, but summer survival is usually poor, especially on upland soils. It produces little fall growth.

Most of the ladino or large varieties flower less than the intermediate varieties and generally depend on their perennial nature for summer survival and production following the seeding year. Osceola is a variety of the ladino type that was

developed by the Florida Agricultural Experiment Station. It flowers more than other ladinos tested in Florida but not as much as LA S-1. It reportedly produces sufficient seed to ensure reseeding in Florida. Its reseeding ability has not been evaluated in Louisiana. Summer survival has been good in south Louisiana tests of Osceola, and yields have generally been higher than for LA S-1.

Several varieties, such as Durana, Regalgraze and Pinnacle, have shown the ability to persist for several years.

Ball Clover: Ball clover is a low-growing winter annual clover. It resembles intermediate white clover. It has excellent reseeding ability. Grazer's Select and Don are currently the only varieties of ball clover available.

Balansa Clover: Balansa clover is a cold-tolerant annual clover that is suited to various soil types. Mature plants are tolerant of periods of saturated soils. It has higher nitrogen production than most clovers, and it is a good reseed. It matures about two weeks later than crimson clover.

Seeding Rates

The seeding rates suggested (Table 4) are based on favorable planting conditions. If soil or climatic conditions are not favorable, the seeding rates should be adjusted upward slightly. Remember that simply planting more seed is not a good substitute for correcting unfavorable conditions that can be corrected.

Many producers plant more than recommended amounts of seed to try to produce more forage or to get earlier grazing. Planting more than the recommended amounts of seed may give slightly earlier grazing in some cases, but the total production for the season is not likely to improve. In planting mixtures of crops, recommended seeding rates must be followed to minimize competition between crops.

Planting Depth

Clovers and other small-seeded legumes should generally be planted at depths of one-quarter to one-half inch. Do not cover them deeper than three-quarters of an inch, or poor stands will result.

Ryegrass and fescue should be planted about one-half inch deep but can tolerate coverage of up to 1 inch under favorable moisture and temperature conditions. They should not be covered deeper than 1 inch. Small grains need soil coverage of about 1 inch for good germination and emergence.

If seedbeds are soft and fluffy, rolling them before planting can help achieve the proper planting depth.

Seeding Dates

Planting pasture and forage crops at the proper date (Table 5) is important to allow good germination and emergence. Planting date can also affect the probability of seedling exposure to insect and disease problems. Planting should be early enough to allow good crop establishment before winter weather becomes severe. The amount of growth produced by plants and the amount of grazing available before winter freezes occur are affected by the planting date. Seed planted into prepared seedbeds can be planted earlier than if planted into undisturbed sod since summer grass sod remains competitive in the early fall planting season. Planting too early into growing sod may result in poor stands or stand failures.

Inoculate Legumes

Forage legumes, when they have the proper rhizobia in root nodules, can provide their own nitrogen and often furnish some additional nitrogen for use by associated grasses. The rhizobia needed are not the same for all legumes and are not even the same for all clovers. Application of the proper rhizobia to the seed before planting (inoculation) is necessary to ensure that they are present. Only packages of rhizobia with labels that specify the legume that is being planted, and for which the expiration date has not passed, should be used. Using an adhesive in applying the inoculant increases the effectiveness of inoculation. Either commercial adhesives or mixtures of water with syrup or sugar can be used. Most clover seed is now commercially coated with several components, including the proper rhizobia inoculant.

| Table 1. Cool-season pasture and forage crop varieties suggested for consideration in 2020-2021. | |
|--|--|
| Crop | Varieties |
| Grasses | |
| Tall Fescue | Jesup Max Q, Kentucky-31 |
| Oats | RAM LA 99016 |
| Cereal Rye | Elbon, Maton, Maton II, Oklon, Wintergrazer 70 |
| Annual Ryegrass | Bashaw Diploid, Bashaw Tetraploid, Diamond T, Double Diamond, Earlyploid, Flying A, FrostProof, Herdsman, Wax Marshall, Nelson Tetraploid, Passerel Plus, Prine, RM4L, TAMTBO, Triangle T and Winterhawk |
| Wheat | (No commercial varieties have been tested in recent years). |
| Triticale | FL 08128 |
| Legumes | |
| Alfalfa | AmeriGraze 702 |
| Arrowleaf Clover | Amclo, Meechi, Yuchi, Apache |
| Balansa Clover | FIXatioN |
| Berseem Clover | Bigbee, Frosty |
| Crimson Clover | Chief, Dixie, Tibbee, AU Robin |
| Red Clover | Kenland, Kenstar, Cherokee, Southern Belle, AU Red Ace, Barduro |
| Subterranean Clover | Mt. Barker, Nangeela, Tallarook, Woogenellup |
| White Clover | LA S-1, Osceola, Regalgraze, Pinnacle, Durana, Neches, Renovation, Cresendo, Stamina |
| Ball Clover | Grazer's Select, Don |

Table 2. Dry forage production (pounds per acre) from small grain entries grown at two locations in Louisiana during three growing seasons (2017-18 through 2019-20).

| | | Location | | |
|-------------------------|-----------|----------------------------|-----------|--------|
| Entry | Species | Franklinton | Winnsboro | Mean |
| | | Dry forage, lb/acre | | |
| 3-Year Production Means | | | | |
| RAM LA99016 | Oat | 4,917 | 3,231 | 4,074 |
| FL 08128 | Triticale | ---- | 2,374 | 2,823† |

†Triticale was evaluated at the Winnsboro location all three years and at Franklinton only in the 2019-2020 growing season so mean over three years represents four year/location environments.



Table 3. Mean dry forage production (pounds per acre) from annual ryegrass entries at three locations in Louisiana during three growing seasons, 2017-2018 through 2019-2020.

| Entry | Location | | | 3-Year Mean |
|-------------------------|-------------------------|------------|-----------|-------------|
| | Franklinton | Jeanerette | Winnsboro | |
| | Dry Forage, lbs. / acre | | | |
| ME-4 (expt.)† | 10,414 | 8,157 | 6,520 | 8,364 |
| Wax Marshall | 10,264 | 8,241 | 6,399 | 8,301 |
| Double Diamond | 10,019 | 8,203 | 5,834 | 8,019 |
| TAMTBO | 10,048 | 8,093 | 5,792 | 7,978 |
| ME-94 (expt.) | 9,078 | 8,589 | 6,218 | 7,962 |
| RM4L | 10,124 | 7,880 | 5,856 | 7,953 |
| Bashaw Tetraploid | 9,268 | 8,240 | 6,224 | 7,911 |
| Prine | 9,758 | 7,984 | 5,983 | 7,908 |
| Herdsmen | 10,097 | 7,950 | 5,557 | 7,868 |
| Diamond T | 9,590 | 8,015 | 5,947 | 7,851 |
| Flying A | 9,011 | 8,588 | 5,891 | 7,830 |
| Deaux Lagniappe (expt.) | 8,949 | 8,448 | 6,014 | 7,804 |
| WMWL (expt.) | 8,894 | 8,428 | 5,956 | 7,759 |
| Tee Lagniappe (expt.) | 8,867 | 7,977 | 6,167 | 7,670 |
| Nelson Tetraploid | 9,375 | 7,568 | 6,038 | 7,661 |
| M2CVS (expt.) | 9,351 | 8,065 | 5,523 | 7,646 |
| Triangle T | 9,440 | 7,579 | 5,894 | 7,638 |
| Winterhawk | 8,999 | 7,873 | 5,921 | 7,597 |
| Bashaw Diploid | 8,685 | 8,122 | 5,923 | 7,577 |
| FrostProof | 8,863 | 7,818 | 5,835 | 7,505 |
| Earlyploid | 9,009 | 7,788 | 5,633 | 7,477 |
| Passerel Plus | 8,357 | 8,359 | 5,703 | 7,473 |
| Gulf (certified) | 8,128 | 7,779 | 5,826 | 7,245 |
| Jackson | 7,867 | 7,800 | 5,913 | 7,193 |
| Mean | 9,269 | 8,064 | 5,940 | 7,758 |
| LSD (.1) | 992 | NS | 395 | 411 |
| CV% | 14 | 10 | 9 | 12 |

†Entries followed by (expt.) are experimental and not commercially available.

Table 4. Seeding rates for cool-season pasture and forage crops.

| Crop | Seeding Rate (lb/A) | |
|---------------------|---------------------|--------------------|
| | Planted Alone | Planted in Mixture |
| Grasses | | |
| Tall Fescue | 30 | 20 |
| Oats | 100 | 60 |
| Cereal Rye | 90 | 50 |
| Annual Ryegrass | 30 | 20 |
| Wheat | 90 | 60 |
| Legumes | | |
| Alfalfa | 20-30 | --- |
| Arrowleaf Clover | 8 | 5 |
| Balansa Clover | 5-8 | 4 |
| Berseem Clover | 20 | 15 |
| Crimson Clover | 15 | 12 |
| Red Clover | 12 | 8 |
| Subterranean Clover | 15 | 12 |
| White Clover | 5 | 3 |
| Ball Clover | 5 | 3 |

Table 5. Planting dates for cool-season pasture and forage crops.

| Crop | Planting Dates for | |
|---------------------|------------------------|-----------------|
| | Prepared Seedbeds | Sod Planting |
| Grasses | | |
| Tall Fescue | Sept. 20-Oct. 15 | --- |
| Oats | Sept. 1-Oct. 15 (N LA) | Approx. Oct. 15 |
| | Sept. 15-Oct.15 (S LA) | Approx. Oct.15 |
| Cereal Rye | Sept. 20-Oct. 15 | Approx. Oct. 15 |
| Annual Ryegrass | Sept. 20-Oct. 15 | Approx. Oct. 15 |
| Wheat | Sept. 20-Oct. 15 | Approx. Oct. 15 |
| Legumes | | |
| Alfalfa | Oct. 5-Oct. 20 | --- |
| Arrowleaf Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Balansa Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Berseem Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Crimson Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Red Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Subterranean Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| White Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |
| Ball Clover | Oct. 1-Nov. 15 | Oct. 15-Nov. 15 |

This material was prepared by the following personnel of the LSU AgCenter:

Dr. Ed Twidwell, Professor, School of Plant, Environmental and Soil Sciences

Dr. M. W. Alison, Associate Professor, Scott Research and Extension Center

Dr. Charles Hutchison, Associate Professor, Southeast Research Station

Greg Williams, Research Associate, Iberia Research Station

Jerry Simmons, Research Associate, Southeast Research Station



Visit our website: www.LSUAgCenter.com

William B. Richardson, LSU Vice President for Agriculture
Louisiana State University Agricultural Center
Louisiana Agricultural Experiment Station
Louisiana Cooperative Extension Service
LSU College of Agriculture

PUB 2334 ONLINE Rev. 8/20

The LSU AgCenter and LSU provide equal opportunities in programs and employment.