

Performance of Grain Sorghum Hybrids in Louisiana, 2010

H.J. “Rick” Mascagni, Jr., Brooks Blanche, Millie Deloach, Bobby Golden, Dustin Harrell, Jim Hayes, Steve Harrison, Clayton Hollier, John Kruse, Roger Leonard, and Boyd Padgett

Performance of grain sorghum hybrids is annually evaluated by Louisiana Agricultural Experiment Station (LAES) researchers. The purpose of these trials is to provide to Louisiana growers, seedsmen, county agents of the Louisiana Cooperative Extension Service (LCES) and other interested individuals and organizations with unbiased performance data for commercial grain sorghum hybrids submitted for evaluation by private agencies.

The cooperating LAES units in 2010 were: Dean Lee Research Station, Alexandria; Central Research Station, Baton Rouge; Red River Research Station, Bossier City; Rice Research Station, Crowley; Northeast Research Station, St. Joseph; and the Macon Ridge Research Station, Winnsboro. Baton Rouge and Crowley locations were lost to very dry conditions.

PROCEDURES

In 2010, 17 grain sorghum hybrids were entered in the LAES yield trials. Soil type, cultural practices, location summaries and weather graphs are listed prior to data tables for each location. In weather graphs, maximum and minimum temperatures are weekly averages and rainfall weekly totals. Trials were not irrigated, except at St. Joseph, where both irrigated and non-irrigated trials were conducted. Seed were treated with Concept and Gaucho and recommended LSU AgCenter cultural practices were followed at each location.

The experimental design at each location was a randomized complete block design with four replications. Traits measured and rating scales are listed in Table 1. Analysis of variance and least significant differences (LSD) were computed using SAS (Statistical Analysis System). We used the protected F-test, which means LSD's were calculated only if differences among hybrids existed at the 90% confidence level. If differences were significant, a LSD at the 10% probability level was calculated. If the LSD (0.10) for yield in

H.J. “Rick” Mascagni, Jr., Professor and Coordinator, Northeast Research Station, St. Joseph, LA 71366; Brooks Blanche, Millie Deloach, and John Kruse Assistant Professor, Research Associate, and Assistant Professor/Specialist, Dean Lee Research Station, Alexandria, LA 71302; Bobby Golden and Jim Hayes, Assistant Professor and Research Associate, Red River Research Station, Bossier City, LA 71113; Steve Harrison, Professor, School of Plant, Environmental and Soil Sciences, Baton Rouge, LA 70803; Clayton Hollier, Professor, Department of Plant Pathology and Crop Physiology, Baton Rouge, LA 70803; Dustin Harrell, Assistant Professor, Rice Research Station, Crowley, LA 70527; and Roger Leonard and Boyd Padgett, Professors, Macon Ridge Research Station, Winnsboro, LA 71295.

a trial is 400 lb/acre, there is a 10% chance that two hybrids with a reported yield difference of 400 lb/acre are genetically equal and a 90% probability they have differences in genetic potential in that particular environment. LSD values are influenced by how well soil fertility, stand establishment, plot length, harvest efficiency and other variables are controlled and by number of replications for each hybrid or treatment. The letters NS are used in the text and tables to indicate lack of significance (**not significantly different**) at the 10% probability level. The coefficient of variation (CV) reflects the magnitude of experimental error (random variation not accounted for by hybrids and replications) in relation to the trial mean. A high CV means that relative differences among hybrids were not consistent among replications, which reduces the precision of a test.

Yields for 2010 and two-year averages (2009 and 2010) are presented in the data tables. Yields for the hybrids in the highest-yielding group for 2010 (yields falling within one LSD value) are in bold print. Hybrids in bold print with a single asterisk fall within the highest-yielding group for both years, 2009 and 2010.

Table 1. Traits and rating scales for LAES grain sorghum performance trials.

Trait	Abbreviation	Description
Yield	Yield	Grain yield @ 14% harvest grain moisture, lb/a
2-Year average yield	2-Year avg	Average grain yield for 2009 and 2010, lb/a
Grain moisture	GrMo	Grain moisture at harvest, %
Test weight	Test wt	Volume weight of grain, lb/bu
Heading date	Mid-head	Date of head emergence in 50% of plants, days after planting (DAP)
Plant height	Plant ht	Plant height from ground to top of head, inches (in)
Head exertion	Head exert	Distance between flag leaf and base of head, inches (in)
Head type	Head type	Head type is a measure of head architecture, with ratings of 1-5; 1-compact, 3-intermediate, and 5-open
Midge damage	Midge	Average percent (%) of head damaged.
Bird damage	Bird	Average percent (%) of head damaged

RESULTS

Yield data and other agronomic data for each location are presented in Tables 2-6. A location summary, soil type, cultural practices and weather information are listed prior to data tables for each location. Yield summary across Louisiana for 2010 is presented in Table 7 and participating seed companies are listed in Table 8.

For additional information on grain sorghum trials, please contact Dr. Rick Mascagni, Northeast Research Station, P.O. Box 438, St. Joseph, LA 71366 (Ph: 318-766-3769; Fax: 318-766-4278; e-mail: hmascagni@agcenter.lsu.edu); or the coordinator at a specific location, Dr. Brooks Blanche, Dean Lee Research Station, Alexandria (Ph: 318-473-6524, Fax 318-473-6535, e:mail: sblanche@agcenter.lsu.edu); Dr. Bobby Golden, Red River Research Station, Bossier City (Ph: 318-741-7430, Fax 318-741-7433, e:mail: bgolden@agcenter.lsu.edu); Dr. Dustin Harrell, Rice Research Station, Crowley (Ph: 337-788-7531, Fax: 337-788-7553, e:mail: dharrell@agcenter.lsu.edu); Dr. Steve Harrison, Central Station, Baton Rouge (Ph:225-578-1308, Fax: 225-578-1403, e:mail:sharrison@agcenter.lsu.edu).